

BISTABILITY FOR ASYMMETRIC DISCRETE RANDOM WALKS

Marco Reyes[*]

Departamento de Física, CINVESTAV, Apdo Postal 14-740, México D.F., México

H. Rosu[*]

Instituto de Física de la Universidad de Guanajuato, Apdo Postal E-143, León, Gto, México

Octavio Obregón[*]

Instituto de Física de la Universidad de Guanajuato, Apdo Postal E-143, León, Gto, México

and Departamento de Física, Universidad Autónoma Metropolitana, Iztapalapa, Apdo Postal 55-534, 09340, Distrito Federal, México

Abstract

We show that asymmetric time-continuous discrete random walks can display bistability for equal values of Jauslin's shifting parameters. The bistability becomes more pronounced at increased asymmetry parameter.

PACS 02.50.Ga - Markov processes

PACS 05.20.Dd - Kinetic theory

In a previous paper [1], two of us presented an extension of a supersymmetric analysis of the three-site discrete master equation due to Jauslin [2]. The essential procedure in this approach is the so-called *addition of eigenvalues (shifting parameters)* method that one may consider as a variant of the negative factorization energy case in the Witten susy scheme as we showed in [1]. Jauslin showed that the supersymmetric master evolution can be put into the following iterative form

$$H_{k-1}\phi_{k-1} = -\lambda_k\phi_{k-1}, \quad k \geq 1 \quad (1)$$

This iterative procedure is based on the initial ($k = 1$) master “Hamiltonian” operator which for homogeneous random walks (jump rates not depending on site) is [1]

$$H_0 = -(c_1 c_2)^{1/2} \frac{\partial^2}{\partial n^2} + c_1 + c_2 - 2(c_1 c_2)^{1/2} \quad (2)$$

c_1 and c_2 are the forward and the backward jump rates, respectively.

Already at the second step ($k = 2$) Jauslin’s method generates bistability in the stationary probability ($H_2 P_2^{st} = 0$) if the second eigenvalue is chosen to be at least two orders of magnitude smaller than the first one as shown in [2] for the case of free random walks ($c_1 = c_2 = 1/2$) and confirmed by us [1]. Multistability can be produced by adding more eigenvalues. This is of course an interesting way of generating bistability and multistability in a master equation.

In this note we want to report on generating bistability in Jauslin’s scheme with the two eigenvalues λ_1 and λ_2 chosen to be equal. We have found that this is possible for asymmetric (or driven) one-step discrete random walks, e.g., possessing jump rates of the type $c_1 = \frac{1}{2}(1 + \epsilon)$ and $c_2 = \frac{1}{2}(1 - \epsilon)$, with ϵ a bias parameter. The idea is to see what happens if we keep the eigenvalues equal (no bistability) but change the other control parameter of the problem which is the degree of asymmetry (or the bias) ϵ . We have found a nice build-up of bistability with increasing ϵ (see the figure) for the stationary state P_2^{st}

$$P_2^{st}(n) = \text{const} \times \frac{2}{(1 - \epsilon^2)^{1/2} \phi_1(n) \phi_1(n-1)} \quad (3)$$

with

$$\phi_1(n) = \frac{(1 - \epsilon^2)^{1/4}}{\sqrt{2}} [(a(n))^{-1/2} \sinh(\gamma_2 n) - (a(n))^{1/2} \sinh(\gamma_2(n-1))] \quad (4)$$

where $a(n) = \frac{\cosh(\gamma_1 n)}{\cosh(\gamma_1(n-1))}$, and $\gamma_1 = \text{arccosh}[\frac{(1+\lambda_1)}{(1-\epsilon^2)^{1/2}}]$, $\gamma_2 = \text{arccosh}[\frac{(1+\lambda_1+\lambda_2)}{(1-\epsilon^2)^{1/2}}]$, with λ_1 and λ_2 being Jauslin’s shifting parameters.

As we said, P_2^{st} is the stationary state of the H_2 “Hamiltonian” which is of the same type as H_0 , i.e.,

$$H_2 = -(c_1'' c_2'')^{1/2} \frac{\partial^2}{\partial n^2} + 1 - 2(c_1'' c_2'')^{1/2} \quad (5)$$

with $c_1'' = \frac{1}{2}(1 - \epsilon^2)^{1/2} \frac{\phi_1(n-1)}{\phi_1(n)}$ and $c_2'' = \frac{1}{2}(1 - \epsilon^2)^{1/2} \frac{\phi_1(n)}{\phi_1(n-1)}$.

A possible application may be to the one-dimensional hopping propagation of charge carriers, such as in (quasi) one-dimensional polymer chains, in the presence of a bias field. In this case the relation between the asymmetry parameter and the electric field is $(1 - \epsilon)/(1 + \epsilon) = \exp(-eEa/kT)$, where E is the bias field, a is the lattice constant, and T is the thermodynamic temperature. One concludes that bistability can show up either in strong fields or at low temperatures.

Acknowledgment

This work was partially supported by the CONACyT Projects 4862-E9406 and 4868-E9406.

M.R. was supported by a CONACyT Graduate Fellowship.

References

- [*] Electronic address: mareyes@fnalv.fnal.gov
- [*] Electronic address: rosu@ifug.ugto.mx
- [*] Electronic address: octavio@ifug.ugto.mx
- [1] H.C. Rosu and M. Reyes, Phys. Rev. E **51**, 5112 (1995)
- [2] H.R. Jauslin, Phys. Rev. A **41**, 3407 (1990)

Figure Caption

Fig. 1 Stationary states $P_2^{st}(n)$ for an asymmetric discrete RW with shifting parameters

$\lambda_1 = \lambda_2 = 0.01$ and asymmetry parameter ϵ as follows:

a) $\epsilon = 0$; b) $\epsilon = 0.25$; c) $\epsilon = 0.50$; d) $\epsilon = 0.75$







Fig. 1